

Supplementary Table 3. Dry-up after three loading injections and recurrence after dry-up, according to the subtypes of nAMD

Variable	Total (n = 38)	Typical nAMD (n = 28)	PCV (n = 10)	<i>p</i> -value
Dry-up after three loading injections				0.462*
Yes	36 (94.7)	27 (96.4)	9 (90.0)	
No	2 (5.3)	1 (3.6)	1 (10.0)	
Recurrence after dry-up (n = 36)				
Yes	28 (77.8)	20 (74.1)	8 (88.9)	0.648*
Recurrence time (mon)	6.5 ± 7.1 (1.9–34.3)	7.2 ± 8.1 (2.0–34.3)	5.0 ± 4.2 (1.9–13.3)	0.238†
No	8 (22.2)	7 (25.9)	1 (11.1)	0.500†
Follow-up period (mon)	29.7 ± 8.1 (22.3–46.9)	30.5 ± 8.4 (22.3–46.9)	24.0	

Values are presented as number (%), mean ± standard deviation (range), or mean.

nAMD = neovascular age-related macular degeneration; PCV = polypoidal choroidal vasculopathy.

*Fisher exact test (typical nAMD vs. PCV); †Mann-Whitney *U*-test (typical nAMD vs. PCV).